

State: HARYANA

Agriculture Contingency Plan District: BHIWANI

1.0 District Agriculture profile				
1.1	Agro-Climatic/Ecological Zone			
	Agro Ecological Sub Region (ICAR)	Western plain hot arid eco-system with desert soils (2.3)		
	Agro-Climatic Region (Planning Commission)	Trans Gangetic Plain region (VI)		
	Agro Climatic Zone (NARP)*	Western Agroclimatic Zone of Haryana (HR-2)		
	List all the districts falling under the NARP Zone	Sirsa, Fatehabad, Hisar, Bhiwani, Mahendragarh, Rewari		
	Geographical coordinates of district	Latitude	Longitude	Altitude
		28° 46' 57.85" N	76° 08' 03.68" E	237 m
	Name and Address of the concerned ZRS/ZARS/RARS/RRTTS	Directorate of Research, CCS HAU, Hissar-125 004		
	Mention the KVK located in the district	Krishi Vigyan Kendra, Near Bhim Stadium, Bhiwani -127 021		
1.2	Rainfall	Average (mm)	Normal Onset	Normal Cessation
	SW monsoon (June-Sep):	340.0	1 st week of July	3 rd week of September
	NE Monsoon(Oct-Dec):	15.7	-	-
	Winter (Jan- March)	36.7		
	Summer (Apr-May)	18.4		
	Annual:	410.8		

* If a district falls in two NARP zone, mention the zone in which more than 50% area falls.

1.3	Land use pattern of the district (latest statistics)	Total geographical area	Forest area	Land under non-agricultural use	Permanent pastures	Cultivable waste land	Land under Misc. tree crops and groves	Barren and uncultivable land	Current fallows	Other fallows
	Area (000 ha)	466	3	24	-	-	-	20	24	-

1.4	Major Soil types	Area ('000 ha)
	Sandy loam soils	139
	Loamy sand soils	167
	Sandy soils	93
	Loamy soils	-
	Clay loam soils	-

1.5	Agricultural land use	Area ('000 ha)	Cropping intensity %
	Net sown area	395	187
	Area sown more than once	345	
	Gross cropped area	740	

1.6	Irrigation	Area ('000 ha)	
	Net irrigated area	283	
	Gross irrigated area	396	
	Rainfed area	112	

Sources of Irrigation	Number	Area ('000 ha)		% area
Canals		144		50.9
Tanks	-	-		-
Open wells	-	-		-
Bore wells	-	139		49.1
Lift irrigation	-	-		-
Other sources	-	-		-
Total	-	283		
Pumpsets	34043	-		-
Micro-irrigation				
Groundwater availability and use	No. of blocks	% area	Quality of water	
Over exploited*	5	56	-	
Critical	-	-	-	
Semi- critical	-	-	-	
Safe	4	44	-	
Wastewater availability and use	-	-	-	
Ground water quality	Alkaline in nature			

*over-exploited: groundwater utilization > 100%; critical: 90-100%; semi-critical: 70-90%; safe: <70%

1.7	Major Field Crops cultivated	Area ('000 ha)*					
		<i>Kharif</i>		<i>Rabi</i>		Summer	Total
		<i>Irrigated</i>	<i>Rainfed</i>	<i>Irrigated</i>	<i>Rainfed</i>		
	Bajra	-	-	--	-	-	184.7
	Rapeseed Mustard	-	-	-	-	-	160.3
	Wheat	-	-	133.2	-	-	133.2

Gram	-	-	-	-	-	69.9
Cotton	48.6	-	-	-	-	48.6
Horticulture crops - Fruits	Total area					
Citrus	0.5					
Ber	0.2					
Guava	0.1					
Horticultural crops - Vegetables	Total area					
Carrot	1.2					
Cauliflower	0.8					
Tomato	0.8					
Medicinal and Aromatic crops	Total area					
Castor	0.2					
Aloe vera	0.01					
Others	0.003					
Plantation crops	-					
Fodder crops	-					
Total fodder crop area	-					
Grazing land	-					
Sericulture etc	-					
Others (Specify)	-					

1.8	Livestock (2008-09)	Male ('000)	Female ('000)	Total ('000)
	Cattle	-	-	94
	Buffaloes total	-	-	472
	Commercial dairy farms	-	-	NA
	Goat	-	-	90
	Sheep	-	-	90
	Others (Camel, Pig, Yak etc)	-	-	39
1.9	Poultry	No. of farms	Total No. of birds ('000)	
	Commercial	NA	818	
	Backyard	NA	4	

1.10	Fisheries						
	A. Capture						
	i) Marine (Data Source: Fisheries Dept.)	No. of fishermen	Boats		Nets		Storage facilities (Ice plants etc.)
			Mechanized	Non-mechanized	Mechanized (Trawl nets, Grill nets)	Non-mechanized (Shore seines, stake & trap nets)	
		-	-	-	-	-	NA
	ii) Inland (Data Source: Fisheries Dept.)	No. Farmer owned ponds		No. of Reservoirs		No. of village tanks	
		NA		NA		NA	
	B. Culture						
		Water Spread Area (ha)		Yield (t/ha)		Production ('000 tons)	
	i) Brakish water (Data source: MPEDA/Fisheries Dept.)	NA		NA		NA	
ii) Fresh water (Data source: Fisheries Dept.)							
Others							

1.11	Production and Productivity of major crops (2007-08)	Kharif		Rabi		Summer		Total		
		Production ('000 t)	Productivity (kg/ha)							
	Bajra	248	1339	-	-			248	1339	
	Rapeseed Mustard	-	-	206	1287	-	-	206	1287	
	Wheat	-	-	527	3964	-	-	527	3964	
	Gram	-	-	56	794	-	-	56	794	
	Cotton	103	357	-	-			103	357	
	Major Horticultural crops									
	Citrus	980	-	-	-	-	-	-	-	980
	Ber	6270	-	--	-	-	-	--	-	6270
	Guava	2010	-	-	-	-	-	-	-	2010

1.12	Sowing window for 5 major crops	Wheat	Cotton	Gram	Bajra	Rapeseed & Mustard
	Kharif- Rainfed	-	-	-	Onset of rain	-
	Kharif-Irrigated	-	15 th April – 7 th July	-	1st week -15th of July	-
	Rabi- Rainfed	October end – November end	-	Mid October – 30 th October	-	September end
	Rabi-Irrigated	October end – 15 th November	-	Mid November – Mid December	-	September end – 20 th October

1.13	What is the major contingency the district is prone to? (Tick mark)	Regular	Occasional	None
	Drought	-	√	-
	Flood	-	-	√
	Cyclone	-	-	√
	Hail storm	-	√	-
	Heat wave	√	-	-
	Cold wave	√	-	-
	Frost	-	√	-
	Sea water inundation	-	-	√
	Pests and diseases (specify)	-	√	-
	Others (Specify)	-	-	-

1.14	Include Digital maps of the district for		
		Location map of district with in State as Annexure I	Enclosed: Yes
		Mean annual rainfall as Annexure 2	Enclosed: Yes
		Soil map as Annexure 3	Enclosed: No

2.0 Strategies for weather related contingencies

2.1 Drought

2.1.1 Rainfed situation

Condition	Major Farming situation	Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Early season drought (delayed onset)	Light textured sandy soils susceptible to wind erosion	Pearl millet: HHB-94, HHB-197, HHB-67 (Improved)	No change	-	-
		Pearl millet + Greengram- Satya, Muskan, Bharpai / Mothbean: RMO 40 (Intercropping 8:4/6:3)	No change	-	
		Clusterbean: HG-563, HG-365 Cowpea: Charodi for grain and CS-88 for fodder Castor: CH-1 Sesame: HT-1 Note- Clusterbean can also inter cropped with pearl millet as above.	No change	-	

Condition	Major Farming situation	Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Early season drought (delayed onset)	Light textured sandy soils susceptible to wind erosion	Pearl millet: HHB-94, HHB-197, HHB-67 (Improved)	Don't grow clusterbean beyond mid July.	-	-
		Pearl millet + Greengram- Satya, Muskan, Bharpai / Mothbean: RMO 40 (Intercropping 8:4/6:3)		-	
		Clusterbean: HG-563, HG-365 Cowpea: Charodi for grain and CS-88 for fodder Castor: CH-1 Sesame: HT-1 Note- Clusterbean can also inter cropped with pearl millet as above.		-	

Condition	Major Farming situation	Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Early season drought (delayed onset)					
Delay by 6 weeks (Aug 3 rd week)	Light textured sandy soils susceptible to wind erosion	Pearl millet: HHB-94, HHB-197, HHB-67 (Improved)	Don't grow sesame beyond mid August.	-	-
		Pearl millet + Greengram- Satya, Muskan, Bharpai / Mothbean: RMO 40 (Intercropping 8:4/6:3)		-	
		Clusterbean: HG-563, HG-365 Cowpea: Charodi for grain and CS-88 for fodder Castor: CH-1 Sesame: HT-1 Note- Clusterbean can also inter cropped with pearl millet as above.		-	

Condition	Major Farming situation	Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Early season drought (delayed onset)					
Delay by 8 weeks (Sept. 1 st week)	Light textured sandy soils susceptible to wind erosion	Pearl millet: HHB-94, HHB-197, HHB-67 (Improved)	Keep fallow	Conserve soil moisture for <i>rabi</i> sowing.	-
		Pearl millet + Greengram- Satya, Muskan, Bharpai / Mothbean: RMO 40 (Intercropping 8:4/6:3)	-do-	-do-	
		Clusterbean: HG-563, HG-365 Cowpea: Charodi for grain and CS-88 for fodder Castor: CH-1 Sesame: HT-1 Note- Clusterbean can also intercropped with pearl millet as above.	-do-	-do-	

Condition	Major Farming situation	Crop/cropping system	Suggested Contingency measures		
			Change in crop/ cropping system	Agronomic measures	Remarks on Implementation
Early season drought (Normal onset)					
Normal onset	Light textured	Pearl millet: HHB-94, HHB-197,	i) In case of poor plant	-	In case of such

followed by 15-20 days dry spell after sowing leading to poor germination/crop stand etc.	sandy soils susceptible to wind erosion	HHB-67 (Improved)	population (<two-third), go for re-sowing as and when rains resume. ii) Gap filling by transplanting under rainy conditions.		situation: i) State Agriculture Department should make arrangement for seeds to meet the exigency at block level. ii) Release of irrigation water in canals and proper power supply may be insured by concerned departments iii) Subsidy on sprinkler, drip irrigation systems and laser leveler
		Pearl millet + Greengram- Satya, Muskan, Bharpai / Mothbean: RMO-40 (Intercropping 8:4/6:3)	In case of poor plant population (<two-third), go for re-sowing as and when rains resume.	-	
		Clusterbean: HG-563, HG-365 Cowpea: Charodi for grain and CS-88 for fodder Castor: CH-1 Sesame: HT-1 Note- Clusterbean can also intercropped with pearl millet as above.	-do-	-	

Condition	Major Farming situation	Crop /cropping system	Suggested Contingency measures		
			Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
Mid season drought (long dry spell, consecutive 2 weeks rainless (>2.5 mm) period)					
At vegetative stage	Light textured sandy soils susceptible to wind erosion	Pearl millet: HHB-94, HHB-197, HHB-67 (Improved)	i) Weeding and hoeing with <i>wheel hand hoe/ kasola</i> as and when required. ii) Thinning to reduce 1/3 rd population.	<i>In-situ/ex-situ</i> moisture conservation: i) Apply life saving irrigation of 4-5 cm, if possible. ii) Foliar spray of urea (2.5 % at 30-35 DAS).	i) Release of irrigation water in canals and proper power supply may be insured by

		Pearl millet + Greengram-Satya, Muskan, Bharpai / Mothbean RMO 40 (Intercropping 8:4/6:3)	i) Don't use chemicals for weed management under stress. ii) Weeding and hoeing with wheel hand hoe/ kasola as and when required. ii) Straw mulching in between rows.	Apply life saving irrigation of 4-5 cm, if possible.	concerned departments ii) subsidy on sprinkler, drip irrigation systems and laser leveler
		Cluster bean: HG-563, HG-365 Cowpea: Charodi for grain and CS-88 for fodder Castor: CH-1 Sesame: HT-1 Note- Clusterbean can also intercropped with pearl millet as above.	-do-	-do-	

Condition	Major Farming situation	Crop /cropping system	Suggested Contingency measures		
			Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
Mid season drought (long dry spell)					
At reproductive stage	Light textured sandy soils susceptible to wind erosion	Pearl millet: HHB-94, HHB-197, HHB-67 (Improved)	i) Remove every third row for green fodder. ii) Make ridge and furrow for rain water harvesting. iii) Life saving irrigation if available.	-	None
		Pearl millet + Greengram- Satya, Muskan, Bharpai / Mothbean: RMO 40 (Intercropping 8:4/6:3)	As above	-	None
		Clusterbean: HG-563, HG-365 Cowpea: Charodi for grain and CS-88 for fodder Castor: CH-1 Sesame: HT-1 Note- Clusterbean can also	As above	-	None

		intercropped with pearl millet as above.			
--	--	--	--	--	--

Condition	Major Farming situation	Crop/cropping system	Suggested Contingency measures		
			Crop management	Rabi crop planning	Remarks on Implementation
Terminal drought (Early withdrawal of monsoon)	Light textured sandy soils susceptible to wind erosion	Pearl millet: HHB-94, HHB-197, HHB-67 (Improved)	Remove every third row for green fodder. Make ridge and furrow for rain water harvesting. Life saving irrigation if available. Foliar spray of urea 2% solution under rainfed condition.	Field preparation for rabi crop sowing during first fortnight of Oct. Sowing of Mustard (RH-30, RH -819, RB-24, RB 50 RH- 781 and Varuna) and Chickpea (C-235, H-208 and HC-1) during second fortnight of Oct.	The State Agriculture Department should have advance arrangements for timely supply of seed, fertilizer and other agro-inputs to farmers at block level. Breeder seed: Dept of Plant Breeding, CCSHAU, Hisar
		Pearl millet + Greengram- Satya, Muskan, Bharpai / Mothbean: RMO 40 (Intercropping 8:4/6:3)	As above	As above	
		Clusterbean: HG-563, HG-365 Cowpea: Charodi for grain and CS-88 for fodder Castor: CH-1 Sesame: HT-1 Note- Clusterbean can also intercropped with pearl millet as above.	As above	As above	

2.1.2 Irrigated situation

Condition	Major Farming situation	Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delayed/ limited release of water in canals due to low rainfall	Sandy soils/sandy loam soils tubewell irrigated	Pearlmillet-Wheat	Pearlmillet-Raya	10-15% higher seed rate Sprinkler irrigation Planting on beds, planting with ridger seeder Laser land leveling, Conjunctive use of canal and ground waters. Intercropping with moong in pearlmillet and harvesting of intercrop Split application of fertilizers Straw mulching Limited ground water use, prefer life saving irrigation Short duration cultivars Soaking of wheat seeds before sowing Seed treatment with azotobactor/rhizobium Deep ploughing during kharif season Shallow irrigation of 4-5 cm depth Weed free environment	Seeds from State, national seed and private seed agencies. The schemes of NREGS, RKRY, NFSM, NHM are in operation. Govt. subsidy on sprinkler, drip irrigation systems and laser leveler
		Pearlmillet-Chickpea	Clusterbean-Barley	10-15% higher seed rate Sprinkler irrigation Planting on beds, planting with ridger seeder Split application of fertilizer Straw mulching Short duration cultivars Seed treatment with azotobactor/rhizobium Deep ploughing during <i>kharif</i> season Shallow irrigation of 4-5 cm depth Weed free environment	
		Fallow -Raya	Summer Greengram-raya	Short duration cultivars Seed treatment with azotobactor/rhizobium Straw mulching Sprinkler irrigation Planting on beds, planting with ridger seeder Laser land leveling,	

				<p>Conjunctive use of canal and ground waters. Marginal ground waters for life saving irrigation Weed free environment</p>	
		Sorghum-Barley	Cucurbits-Raya	<p>Sprinkler irrigation Planting on beds, planting with ridger seeder Laser land leveling, Conjunctive use of canal and ground waters. Split application of fertilizer Straw mulching Limited ground water use, prefer life saving irrigation Seed treatment with azotobactor Deep ploughing during kharif season Shallow irrigation of 4-5 cm depth Weed free environment</p>	
	Well drained, medium alluvial soils, canal and tubewell irrigated	Clusterbean-Wheat	Cotton-wheat	<p>Drip/furrow irrigation in cotton, paired row planting Sprinkler in wheat Planting on beds Straw mulching in cotton Planting on beds Planting with ridger seeder Laser land leveling Split application of fertilizer Straw mulching in sugarcane Limited ground water use, prefer life saving irrigation Conjunctive use of brackish ground waters with canal waters Short duration cultivars Soaking of wheat seeds before sowing Seed treatment with azotobactor/rhizobium Deep ploughing during <i>kharif</i> season Shallow irrigation of 4-5 cm depth Sowing of vegetable seeds in polythene bags and replanting them in holes. Weed free environment</p>	<p>Shallow ground water use alone or in combination. Seeds from State, national and private seed agencies seed agencies, The schemes of NREGS, RKRY, NFSM, NHM are in operation. Govt. subsidy on sprinkler and drip irrigation systems, on laser land leveling</p>
		Pearlmillet-Wheat	Pearlmillet-Raya/chickpea	<p>Paired row planting Sprinkler irrigation Planting on beds Straw mulching Laser land leveling Split application of fertilizer</p>	

				<p>Straw mulching Limited ground water use, prefer life saving irrigation Conjunctive use of brackish ground waters with canal waters Short duration cultivars Seed treatment with azotobactor/rhizobium Deep ploughing during <i>kharif</i> season Shallow irrigation of 4-5 cm depth Weed free environment</p>	
		Sugarcane-wheat	Sugarcane– Greengram intercropping	<p>Drip/furrow irrigation in sugarcane, paired row planting Planting on beds Straw mulching in sugarcane Laser land leveling Split application of fertilizer Limited ground water use, prefer life saving irrigation Conjunctive use of brackish ground waters with canal waters Short duration cultivars Weed free environment Weed free environment</p>	
		Pearlmillet/fallow- raya	Vegetables	<p>Furrow irrigation in pearlmillet/raya, paired row planting Planting on beds Straw mulching Laser land leveling Split application of fertilizer Marginal ground waters for life saving irrigation Conjunctive use of brackish ground waters with canal waters Short duration cultivars Seed treatment with azotobactor Deep ploughing during <i>kharif</i> season Shallow irrigation of 4-5 cm depth Sowing of vegetable seeds in polythene bags and replanting them in holes. Weed free environment</p>	
	Clay soils, canal and tubewell	Rice-wheat	Summer Greengram-Rice	<p>Sprinkler irrigation in moong, Planting on beds</p>	<p>Late sown cultivars Short duration Desi wheat</p>

	irrigated			Laser land leveling	and Basmati rice. Shallow ground water use alone or in combination. Conservation of rain water, mulching, rain water harvesting Seeds from State and national seed agencies, The schemes of NREGS, RKRY, NFSM, NHM are in operation. Seed from private seed agencies
		Sugarcane-wheat	Sugarcane-onion intercropping	Drip irrigation in paired row planting of sugarcane Laser land leveling Straw mulching in sugarcane	
		Sorghum fodder-wheat	Vegetables/ Flowers	Sprinkler/drip irrigation, Planting on beds, laser land leveling Mulching on inter-row spacing Use of marginal ground waters as life saving irrigation	

Condition	Suggested Contingency measures				Remarks on Implementation
	Major Farming situation	Crop/cropping system	Change in crop/cropping system	Agronomic measures	
Non release of water in canals under delayed onset of monsoon in catchment	Sandy soils, tubewell irrigated	Pearlmillet-Raya	Pulses-Raya	Planting on beds Sprinkler irrigation Marginal ground waters for life saving irrigation Laser land leveling Straw mulching Paired row planting Split application of fertilizer Straw mulching Marginal ground waters for life saving irrigation Conjunctive use of brackish ground waters with canal waters Short duration cultivars Seed treatment with azotobactor/rhizobium Deep ploughing during <i>kharif</i> season Shallow irrigation of 4-5 cm depth Weed free environment	Short duration cultivars of crops Shallow ground water use alone or in combination. Conservation of rain water, mulching, rain water harvesting. Shallow ground water use alone or in combination. Conservation of rain water, mulching, rain water harvesting.
		Pearlmillet-Chickpea	Clusterbean-Barley	Sprinkler irrigation Planting on beds Straw mulching	

				<p>Laser land leveling Split application of fertilizer Marginal ground waters for life saving irrigation Conjunctive use of brackish ground waters with canal waters Short duration cultivars Seed treatment with azotobactor Deep ploughing during kharif season Shallow irrigation of 4-5 cm depth Weed free environment</p>	
		Fallow – Raya/Barley	Vegetables-Raya	<p>Sowing of vegetable seeds in polythene bags and replanting them in holes. Drip irrigation in vegetables Planting on beds Straw mulching Laser land leveling Split application of fertilizer Limited ground water use, prefer life saving irrigation Conjunctive use of brackish ground waters with canal waters Seed treatment with azotobactor Deep ploughing during <i>kharif</i> season Shallow irrigation of 4-5 cm depth Weed free environment</p>	
	Well drained, medium alluvial soils, canal and tubewell irrigated	Clusterbean- Barley	Cotton-Wheat	<p>Drip/furrow irrigation in cotton Sprinkler in wheat Planting on beds Laser land leveling Limited ground water use, prefer life saving irrigation Conjunctive use of ground water Shallow irrigation of 4-5 cm depth Weed free environment</p>	<p>Short duration cultivars of crops Shallow ground water use alone or in combination. Conservation of rain water, mulching, rain water harvesting. Shallow ground water use alone or in combination. Conservation of rain water, mulching, rain water harvesting</p>
		Pearlmillet/fallow- Wheat	Pearlmillet- Raya/Chickpea	<p>Paired row planting Sprinkler irrigation Planting on beds Straw mulching Laser land leveling</p>	

				<p>Split application of fertilizer Straw mulching Limited ground water use, prefer life saving irrigation Conjunctive use of brackish ground waters with canal waters Short duration cultivars Seed treatment with azotobactor/rhizobium Deep ploughing during kharif season Shallow irrigation of 4-5 cm depth Weed free environment</p>	
		Pearlmillet/fallow-Raya	Sugarcane+Greengram intercropping	<p>Drip/furrow irrigation in sugarcane, paired row planting Planting on beds Straw mulching in sugarcane Laser land leveling Split application of fertilizer Limited ground water use, prefer life saving irrigation Conjunctive use of brackish ground waters with canal waters Short duration cultivars Weed free environment</p>	
		Sorghum -Wheat	Vegetables	<p>Sowing of vegetable seeds in polythene bags and replanting them in holes. Drip irrigation in vegetables Planting on beds Straw mulching Laser land leveling Split application of fertilizer Limited ground water use, prefer life saving irrigation Conjunctive use of brackish ground waters with canal waters Seed treatment with azotobactor Deep ploughing during kharif season Shallow irrigation of 4-5 cm depth Weed free environment</p>	
	Clay soils, canal	Pigeon pea –	Summer Moong-		Short duration cultivars of

	and tubewell irrigated	Wheat/barley Fallow --Raya	Wheat Sugarcane +Mungbean intercropping	Drip/furrow irrigation in sugarcane, paired row planting Planting on beds Straw mulching in sugarcane Laser land leveling Split application of fertilizer Limited ground water use, prefer life saving irrigation Conjunctive use of brackish ground waters with canal waters Short duration cultivars Weed free environment	crops Shallow ground water use alone or in combination. Conservation of rain water, mulching, rain water harvesting.
		Sorghum fodder- Wheat	Vegetables/ flowers	Sowing of vegetable seeds in polythene bags and replanting them in holes. Drip irrigation in vegetables Planting on beds Straw mulching Laser land leveling Split application of fertilizer Limited ground water use, prefer life saving irrigation Conjunctive use of brackish ground waters with canal waters Seed treatment with azotobactor /rhizobium	

Condition	Major Farming situation	Crop/cropping system	Change in crop/cropping system	Suggested Contingency measures	
				Agronomic measures	Remarks on Implementation
Lack of inflows into tanks due to insufficient /delayed onset of monsoon	Sandy soils, tubewell irrigated	Pearlmillet-Wheat	Clusterbean-Wheat	Planting on beds, sprinkler irrigation, Planting on beds, sprinkler irrigation, Drip irrigation Limited ground water use, prefer life saving irrigation	Short duration cultivars of crops Shallow ground water use alone or in combination. Conservation of rain water, mulching, rain water harvesting. Shallow ground water use
		Sorghum-Wheat	Sugarcane-Wheat/raya		
		Pearlmillet-Chickpea	Fallow-Raya		

					alone or in combination. Conservation of rain water, mulching, rain water harvesting
Well drained, medium alluvial soils, canal and tubewell irrigated	Rice-Wheat	Pearlmillet-Chickpea	Drip/furrow irrigation in cotton, sprinkler in wheat, planting on beds Sprinkler irrigation, Planting on beds, planting with ridger seeder, laser land leveling Limited ground water use, prefer life saving irrigation Drip irrigation, paired row Planting Drip irrigation, paired row planting Sprinkler irrigation, Planting on beds, planting with ridger seeder, laser land	-do-	
	Sugarcane-Wheat	Pigeonpea-Wheat			
	Rice- Berseem(fodder)	Cotton-Wheat			
Clay soils, canal and tubewell irrigated	Pigeonpea – Wheat/barley	Summer Greengram- Wheat	Drip irrigation, paired row planting of sugarcane Planting on beds Shallow irrigation in vegetable and straw mulching Conjunctive use of ground water Use of gypsum for reclaiming sodic waters Limited ground water use, prefer life saving irrigation	-do-	
	Sugarcane-Wheat	Sugarcane-Greengram intercropping			
	Sorghum fodder- Wheat	Vegetables/ flowers			

Condition	Major Farming situation	Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Insufficient groundwater recharge due to low rainfall	Sandy soils, tubewell irrigated	Pearlmillet-Barley	Clusterbean-Wheat	Adoption of efficient methods of irrigation viz., drip in wide spaced, vegetables and horticultural crops Sprinkler irrigation in other crops	Artificial ground water recharge
		Fallow-Raya	Sugarcane-Wheat/Raya		
		Pearlmillet-Chickpea	Fallow-Raya		
	Well drained, medium alluvial soils, canal and tubewell irrigated	Rice-Wheat	Pearlmillet-Chickpea		
		Sugarcane-Wheat	Pigeonpea-Wheat		
		Rice-Berseem (fodder)	Cotton-Wheat		
	Clay soils, canal and	Pigeon pea –Wheat/Barley	Clusterbean-Raya		

	tubewell irrigated	Pearlmillet–Raya/Chickpea	Planting on beds		
		Sorghum fodder-Wheat	Cucurbits-Raya		

2.2 Unusual rains (untimely, unseasonal etc)

Condition	Suggested contingency measure			
	Vegetative stage	Flowering stage	Crop maturity stage	Post harvest
Continuous high rainfall in a short span leading to water logging				
Rice	Drainage, if depth of standing water is > 5-6 cm	Drainage	Drainage	Shifting to dry place
Cotton	Drainage	-do-	-do-	-do-
Pearlmillet	-do-	-do-	-do-	-do-
Sorghum (fodder)	-do-	-do-	-do-	-do-
Horticulture				
All crops	<ol style="list-style-type: none"> 1. No adverse effect 2. Removal of unwanted sprouts 3. Spray insecticides & pesticides to control the insect & pest 4. Drain out water if heavy rains 	<ol style="list-style-type: none"> 1. Drain out the excess water to avoid flower and fruit drop 2. To control the fruit drop apply foliar application of nutrients and growth regulators 3. Apply insecticide & pesticides to control the insect & pest and diseases on young developing fruits 4. Plough the field to increase the root aeration. 	Harvest the fruit crops timely and send to the market immediately.	<ol style="list-style-type: none"> 1. Apply fungicide to avoid post harvest diseases. 2. Proper covering of the produce. 3. Proper grading and cleaning of fruits immediately after harvest. 4. Use the damaged fruits for processing 5. Use water proof packaging
Heavy rainfall with high speed winds in a short span				
Rice	Drainage, if stagnant water	Drainage	Drainage	Shifting to dry place
Cotton	-do-	-do-	-do-	-do-
Pearlmillet	-do-	-do-	-do-	-do-
Sorghum (fodder)	-do-	-do-	-do-	-do-
Horticulture				

All crops	<ol style="list-style-type: none"> 1. No adverse effect 2. Removal of unwanted sprouts 3. Spray insecticides & pesticides to control the insect & pest 4. Drain out water if heavy rains 	<ol style="list-style-type: none"> 1. Drain out the excess water to avoid flower and fruit drop 2. To control the fruit drop apply foliar application of nutrients and growth regulators 3. Apply insecticide & pesticides to control the insect & pest and diseases on young developing fruits 4. Plough the field to increase the root aeration. 	Harvest the fruits and send to the market immediately.	<ol style="list-style-type: none"> 1. Apply fungicide to avoid post harvest diseases. 2. Proper covering of the produce. 3. Proper grading and cleaning of fruits immediately after harvest. 4. Use the damaged fruits for processing 5. Use water proof packaging
Outbreak of pests and diseases due to unseasonal rains				
Wheat : Yellow and brown rust of wheat become severe Karnal bunt infection increases under moist conditions	Spray 600 – 800 gm Mancozeb 200 lt. of water/acre at the appearance of disease and repeat after 15-20 days			Treat wheat seed with Raxil 2DS @ 1 gm/kg before sowing to control Karnal bunt
Bajra : Downy mildew incidence increases	There is no control measure except resistant varieties			
Indian Mustard: White rust and Alternaria leaf blight increase, stem rot increases due to rain and cold weather	Spray Mancozeb 0.2% 3-4 times at an interval of 15 days to control white rust and Alternaria leaf blight.	To control stem rot spray 0.2% Carbendazim.		
Cotton : Bacterial leaf blight increases due to rainfall from traces to moderate intensity whereas cotton leaf curl virus decreases	Soak 5 -6 kg delimited and limited cotton seed in 10 lt. of water suspension containing 5 g Emisan + 1 gm Streptocycline sulphate for 2 hrs. and 6-8 hrs respectively before sowing..			
Horticulture				
Potato: Early blight of potato increases with rainfall	Spray Mancozeb @ 0.25% 4-5 times at an interval of 15 days			

2.3 Floods

Condition	Suggested contingency measure			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Transient water logging/ partial inundation				
Rice	Surface drainage	Drainage	Drainage	Shifting to dry place
Cotton	-do-	-do-	-do-	-do-
Pearlmillet	-do-	-do-	-do-	-do-
Sorghum	-do-	-do-	-do-	-do-
Horticulture				
All crops	<ul style="list-style-type: none"> ➤ Drain out the flood water ➤ Spray of nutrients/supplementation ➤ Prefer plantation of water logging resistant crop like Jamun. ➤ Mount planting of fruit trees 			Drain out the flood water
Continuous submergence for more than 2 days				
Rice	Surface drainage	Drainage	Drainage	Shifting to dry place
Cotton	-do-	-do-	-do-	-do-
Pearlmillet	-do-	-do-	-do-	-do-
Sorghum	-do-	-do-	-do-	-do-
Horticulture				
All crops	<ul style="list-style-type: none"> ➤ Drain out the flood water ➤ Spray of nutrients/supplementation ➤ Prefer plantation of water logging resistant crop like Jamun. ➤ Mount planting of fruit trees 			Drain out the flood water
Sea water inundation	NA			

2.4 Extreme events: Heat wave / Cold wave/Frost/ Hailstorm /Cyclone

Extreme event type	Suggested contingency measure			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Heat Wave				
Rice	Micro-irrigation, avoid irrigation during hot hours with poor quality waters	Micro-irrigation, avoid irrigation during hot hours with poor quality waters	-	
Cotton	Micro-drip irrigation	Deep irrigation	Deep irrigation	
Pearlmillet	Micro-sprinkler irrigation, avoid irrigation during hot hours with poor quality waters	Micro- sprinkler irrigation, avoid irrigation during hot hours with poor quality waters	Micro-irrigation, avoid irrigation during hot hours with poor quality waters	
Sorghum	-do-	-do-	-do-	
Clusterbean	-do-	-do-	-do-	

Extreme event type	Suggested contingency measure			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Pigeonpea	-do-	-do-	-do-	
Horticulture				
All crops	Micro-irrigation, avoid irrigation during hot hours with poor quality waters	-do-	-do-	
Cold wave				
Wheat	Irrigation, balanced fertilizer application, Foliar spray of nutrients	Irrigation, fertilizer application	Irrigation, fertilizer application	
Raya	-do-	-do-	-do-	
Chickpea	-do-	-do-	-do-	
Barley	-do-	-do-	-do-	
Fodder	-do-	-do-	-do-	
Horticulture				
Frost				
Wheat	No adverse effect			
Raya	Irrigate the crop Create smoke during late evening	Irrigate the crop Create smoke during late evening	Irrigate the crop Create smoke during late evening	
Chickpea	-do-	-do-	-do-	
Barley	-do-	-do-	-do-	
Fodder	-do-	-do-	-do-	
Horticulture				
All crops	<ol style="list-style-type: none"> 1. Apply light irrigation frequently 2. Creating smoke in the orchard during late evening. 3. Thatching of young plants during severe cold months. 4. Use of sprinkler irrigation. 5. Use of mulching under plant canopy 			
Hailstorm				
Horticulture	Use of Anti-hail nets			
	<ol style="list-style-type: none"> i. Plantation of wind breakers ii. Use of hailstorm nets iii. Supplementation of nutrients to the trees 			
Cyclone				
	-			
Horticulture				
All crops	Seedling covers should be used			

2.5 Contingent strategies for Livestock, Poultry & Fisheries

2.5.1 Livestock

	Suggested contingency measures		
	Before the event	During the event	After the event
Drought			
Feed and fodder availability	<ol style="list-style-type: none"> All Districts should be asked to locate their feed and fodder banks in view of submergence situation arising due to draught. Sufficient care must be taken to sensitize the farmers to protect their feed and fodder much ahead of onset of monsoon. The sources for procurement of feed / rice bran (Kunda) within the district and nearest locations should be identified, and the suppliers kept informed about the emergency situation, which might require action at their level for production and supply to the identified areas within the shortest possible time. Complete feed blocks should be prepared and stored in the feed banks for scarcity periods. The livestock holders of small ruminants should be educated/ informed to collect sufficient amount of green leaves from edible plants for use during the period of submergence at the earliest, after receipt of draught warning. The district authorities of Animal Husbandry Department should chalk out a complete programme to cater the feed & fodder needs of livestock. Increase the sown area under fodder crops Looking to scarcity of crop residues, burning of paddy straw and stubbles should not be allowed in Haryana. This can be properly harvested, baled, densified and fortified using 4% urea with molasses and transported to areas of fodder scarcity. Standardized machinery for harvesting, bailing, densification and fortification is available with 	<ol style="list-style-type: none"> The best option is to open fodder depots for milch animals which farmers will never deposit into the cattle camps and establish cattle camps for dry and scrub animals. These camps should be established along assured source of water or canals for drinking and growing fodder. Facilities like storing densified roughages transported from other districts should also be established adjacent to these camps. Complete feed blocks stored in the feed banks should be provided to productive, lactating and pregnant animals for scarcity periods Since stall feeding adversely affects the breeding efficiency in case of sheep, therefore, sheep should always be resorted to natural grazing. Special care is required for productive, lactating and pregnant animals. These animals must be supplemented with additional concentrates and fodders. Most of such animals will be retained by the farmers and arrangements for fodder, feed and drinking water should be made accordingly. 	<ol style="list-style-type: none"> Immediate efforts are needed to grow short duration fodder crops like oats, barley, <i>kasni</i> and <i>lucern</i> etc. in the canal command areas. Farmers might have to be compensated for abandoning food or commercial cash crop to meet contingent fodder requirements.

	Suggested contingency measures		
	Before the event	During the event	After the event
	Punjab Agro Federation and in the market.		
Drinking water	Prior to the onset of summer all the water ponds/lakes in the villages/cities should be filled up with canal water/tube wells.	<ol style="list-style-type: none"> 1. All the affected livestock should have an access to clean drinking water. Arrangements are required to be made in this regard with the help of concerned Government functionaries of the Districts. 2. Resorting to alternate day watering to camel, sheep and goats. Experimental evidences show that even watering twice a week did not have much adverse effect on body weight of the sheep. 3. Avoiding long distance grazing, as tired animals need more and frequent watering and feeding. 	Normal supply of water should be restored.
Health and disease management	Constitution of task force at district and sub division level which will formulate guidelines for action should have a mobile veterinary unit at their disposal. Procurement of mineral and feed supplements, life saving drugs, electrolytes, vaccines etc.	Disbursement of supplements, treatment of affected animals in camps, proper disposal of dead animals, deworming and vaccinations.	Rehabilitation of affected animals, provision of veterinary aid and follow up, provide supplements etc to make up losses for deficiencies.
Floods			
Feed and fodder availability	<ol style="list-style-type: none"> 1. All Districts should be asked to locate their feed and fodder banks in view of submergence situation arising due to floods. Sufficient care must be taken to sensitize the farmers to protect their feed and fodder much ahead of onset of monsoon. The sources for procurement of feed / rice bran (Kunda) within the district and nearest locations should be identified, and the suppliers kept informed about the emergency situation, which might require action at their level for production and supply to the identified areas within the shortest possible time. 2. Complete feed blocks should be prepared and stored in the feed banks for scarcity periods 3. The livestock holders of small ruminants should be educated/ informed to collect sufficient amount of green leaves from edible plants for use during the period of submergence at the earliest, after receipt 	<ol style="list-style-type: none"> 1. The best option is to open fodder depots for milch animals which farmers will never deposit into the cattle camps and establish cattle camps for dry and scrub animals. These camps should be established along assured source of water or canals for drinking and growing fodder. 2. Facilities like storing densified roughages transported from other parts of the country should also be established adjacent to these camps. 3. Immediate efforts are needed to grow fodder crops like oats, barley, <i>kasni</i> and <i>lucern</i> etc. in the canal command areas. 4. Farmers might have to be compensated for abandoning food or commercial cash crops to meet contingent fodder requirements. 5. Since stall feeding adversely affects the breeding efficiency in case of sheep, therefore, sheep should 	<ol style="list-style-type: none"> 1. Immediate efforts are needed to grow fodder crops like oats, barley, <i>kasni</i> and <i>lucern</i> etc. in the canal command areas. 2. Farmers might have to be compensated for abandoning food or commercial cash crops to meet contingent fodder requirements. 3. After the sheds have dried, these should be disinfected and regular feed of the animals should be introduced gradually.

	Suggested contingency measures		
	Before the event	During the event	After the event
	<p>of draught warning. The district authorities of Animal Husbandry Department chalk out a complete programme to cater the feed & fodder needs of cattle, buffalo, sheep, goat, pig, dog, poultry birds etc.</p> <p>4. The livestock holders of livestock are trained regarding shifting of animals before flooding. The farmers are instructed to let loose their animals instead of tying much before flood.</p> <p>5. Increase the sown area under fodder crops</p> <p>6. Looking to scarcity of crop residues, burning of paddy straw and stubbles should not be allowed in Haryana. This can be properly harvested, baled, densified and fortified using 4% urea with molasses and transported to areas of fodder scarcity. Standardized machinery for harvesting, baling, densification and fortification is available with Punjab Agro Federation and in the market.</p>	<p>always be resorted to natural grazing.</p> <p>6. Special care is required for productive, lactating and pregnant animals. These animals must be supplemented with additional concentrates and fodders.</p> <p>7. Most of such animals will be retained by the farmers and arrangements for fodder, feed and drinking water should be made accordingly.</p>	
Drinking water	<p>Tube wells should be installed before monsoon to provide underground water to the livestock during flood period.</p>	<p>All the affected livestock and poultry should have an access to clean drinking water. Arrangements are required to be made in this regard with the help of concerned Government functionaries of the Districts. The available water may be chlorinated if required with help of Halogen Tablet prior to drinking by livestock and poultry.</p>	<p>Normal supply of water should be restored.</p>
Health and disease management	<p>Constitution of task force at district and sub division level which will formulate guidelines for action. Procurement of mineral and feed supplements, life saving drugs, electrolytes, vaccines etc. Workout places for evacuation.</p>	<p>Evacuate to safe places, provide veterinary aid to affected animals, proper disposal of dead animals, disinfection of drinking water. If not already done, carry out deworming and vaccinations for HS, FMD, BQ in cattle, PPR, sheep pox, ET in sheep and goats, swine fever in pigs..</p>	<p>Rehabilitation of affected animals, provision of veterinary aid and follow up, provide supplements etc. Disinfection of area, control of vectors, prevention of spread of disease/outbreaks. Treatment of affected animals.</p>
Cyclone	-NA-		
Feed and fodder availability			

	Suggested contingency measures		
	Before the event	During the event	After the event
Drinking water			
Health and disease management			
Heat wave and cold wave			
Shelter/environment management	Necessary arrangement of tatties, gunny bags and tirpal should be made available so as to cover the sheds during heat and cold waves	1. Window of the sheds should be covered with gunny bags, tatties, and tirpal. Electric fans should be provided in the sheds and if possible desert cooler should be provided during heat period. 2. High energy and readily available sources of energy nutrients may be provided in the ration.	Normal shelter should be restored
Health and disease management	Provision of shelter/roof/covered and open area to animals, procurement of life saving drugs and vaccines.	Cold waves: Cover the animal with old blanket/gunny bag etc. Heat wave: Sprinkle water/take buffaloes to ponds. Treat affected animals, vaccinate if not done earlier.	Treatment of affected animals, provide veterinary aid and follow up.

2.5.2 Poultry

	Suggested contingency measures		
	Before the event	During the event	After the event
Drought			
Shortage of feed ingredients	I. All Districts should be asked to locate their feed banks in view of submergence situation arising due to draught. Sufficient care must be taken to sensitize the farmers to protect their feed and fodder much ahead of onset of monsoon. The sources for procurement of feed / rice bran (Kunda) within the district and nearest locations should be identified, and the suppliers kept informed about the emergency situation, which might require action at their level for production and supply to the identified areas within the shortest possible time. II. The district authorities of Animal Husbandry Department should chalk out a complete programme to cater to feed the poultry birds.	Poultry farmers should be provided with sufficient amount of feed ingredients and complete feed during draught situation from the feed banks.	Normal feeding should be restored
Drinking water	Necessary arrangement for water storage should be made. Hand pumps should be installed around the sheds. Sufficient quantity of electrolytes	All the affected poultry should have an access to clean drinking water. Arrangements are	Normal drinking water restored

	should be ensured.	required to be made in this regard with the help of concerned Government functionaries of the Districts.	
Health and disease management	Constitution of task force at district and sub division level which will formulate guidelines for action should have a mobile veterinary unit at their disposal. Commercial poultry farms can procure grain/feed in advance.	In backyard birds, put some grains and sufficient water inside the enclosure, provide some vitamin supplement.	In backyard poultry, carry out de-worming and vaccination for Ranikhet disease and Gumboro. Provide vitamins and mineral supplement.
Floods			
Shortage of feed ingredients	<p>I. All Districts should be asked to locate their feed banks in view of submergence situation arising due to flood. Sufficient care must be taken to sensitize the farmers to protect their feed much ahead of onset of monsoon. The sources for procurement of feed / rice bran (Kunda) within the district and nearest locations should be identified, and the suppliers kept informed about the emergency situation, which might require action at their level for production and supply to the identified areas within the shortest possible time.</p> <p>II. The poultry farmers should be trained regarding shifting of birds before flood. For shifting of poultry birds to safer places, the farmer should be educated to make suitable cages from bamboos.</p>	Sufficient quantity of feeds stored in the feed banks should be made available to the poultry farmers.	Normal feeding should be restored
Drinking water	I. Prior to the onset of monsoon tube wells should be installed in the villages and near to the poultry farms so as to provide underground water during flood.	All the affected poultry should have an access to clean drinking water. Arrangements are required to be made in this regard with the help of concerned Government functionaries of the Districts. The available water may be chlorinated if required with help of Halogen Tablet prior to drinking by livestock and poultry.	Normal drinking water restored
Health and disease management	Constitution of task force at district and sub division level which will formulate guidelines for action should have a mobile veterinary unit at their disposal. Make provision of shelter for evacuation and arrangement around farm so that flood water does not enter poultry farm/shed. Provision or facilities for disposal of dead birds.	Evacuate the birds to safer places. Carry out deworming and vaccinations. May dispose off/sell birds for meat purpose. Proper disposal of dead birds.	Make shed dry, sprinkle lime & spray insecticides, disinfectant before placement of birds,

			use of coccidiostat in feed or water, proper disposal of dead birds.
Cyclone	-NA-		
Shortage of feed ingredients			
Drinking water			
Health and disease management	Keep arrangements in place in shed for heating during winter/cold waves and for cooling by use of sprinklers/foggers. Procure electrolytes and supplements.	Avoid too much fluctuation below the temperature of 70 °F and above 100 °F. Use bukharies, gas burner, secure curtains during winter. Provide a course of antibiotics in feed or water for 3-5 days to combat respiratory problems. Provide vitamin C, electrolyte in drinking water during heat waves and use of foggers, wetting of curtains, sprinkling of water etc. during heat waves. May dispose off/sell birds if heavy mortality occurring.	Treatment of affected birds, vaccination if delayed may be carried out as per schedule.
Heat wave and cold wave			
Shelter/environment management	Necessary arrangement of <i>tatties</i> , gunny bags and <i>tirpal</i> should be made available so as to cover the sheds during heat and cold waves	Window of sheds should be covered with gunny bags, tatties, & tirpal. Electric fans should be provided in the sheds and if possible desert cooler should be provided during heat period. High energy & readily available sources of energy nutrients may be provided in ration.	Normal shelter should be restored
Health and disease management			

2.5.3 Fisheries

	Suggested contingency measures		
	Before the event	During the event	After the event
1) Drought			
A. Capture	NA		

Marine			
Inland			
(i) Shallow water depth due to insufficient rains/inflow			
(ii) Changes in water quality			
B. Aquaculture			
(i) Shallow water in ponds due to insufficient rains/inflow	Further increase the depth of ponds, store the fish stock in 1 & 2 ponds only.	Sell the big fishes and keep the smaller fishes in one tank.	Stock the young fishes in different tanks, species wise.
(ii) Impact of salt load build up in ponds / change in water quality	Continuously add some water from tube well/water source in fish ponds	Do not allow the water level to go below 3.5 feet in fish ponds.	Stock the young fishes in different tanks and keep the water between 3.5 and 6.0 feet.
2) Floods	NA		
A. Capture			
Marine			
Inland			
(i) No. of boats / nets/damaged			
(ii) No.of houses damaged			
(iii) Loss of stock			
(iv) Changes in water quality			
(v) Health and diseases			
B. Aquaculture			
(i) Inundation with flood water	Boundaries/bunds with height >6 feet may be made around fish ponds, will restrict, escape of fishes from ponds	Net-out and stock the fishes in one big tanks and make the bund >6 feet height around the ponds.	Remove the bund separately and release the fishes, species-wise in tanks.
(ii) Water contamination and changes in water quality	Add more fresh water in each tank (tube well/canal), grow aquatic weeds.	Repeatedly filter and re-circulate water from stocking tanks	Filter, re-circulate and add new fresh water every week, will decrease fish mortality.

(iii) Health and diseases	Treat the pond water with KmNO_4 @ 10 ppm in each fish tanks. Add new fresh water periodically.	Disinfect fish ponds with KmNO_4 @ 10g/10,000 liter water fortnightly.	Treatment with KmNO_4 must continue for one month even after flood situation is out. Remove the highly infected fishes from ponds.
(iv) Loss of stock and inputs (feed, chemicals etc)	Store the inputs at safer places.	Move stock and inputs to safer places and acquire fresh stock in shortage.	Retain the normal arrangements.
(v) Infrastructure damage (pumps, aerators, huts etc)	Make alternate arrangements according to the anticipated conditions	Proper maintenance/repairing of damaged infrastructure or make new arrangements.	Proper maintenance/repairing of damaged infrastructure.
3. Cyclone / Tsunami	NA		
A. Capture			
Marine			
(i) Average compensation paid due to loss of fishermen lives			
(ii) Avg. no. of boats / nets/damaged			
(iii) Avg. no. of houses damaged			
Inland			
B. Aquaculture			
(i) Overflow / flooding of ponds			
(ii) Changes in water quality (fresh water / brackish water ratio)			
(iii) Health and diseases			
(iv) Loss of stock and inputs (feed, chemicals etc)			
(v) Infrastructure damage (pumps, aerators, shelters/huts etc)			
4. Heat wave and cold wave			

A. Capture	NA		
Marine			
Inland			
B. Aquaculture			
(i) Changes in pond environment (water quality)	Keep the ponds water fresh by adding fresh tubewell water, regularly.	Showering the water in air and add fresh tube-well water, periodically.	During heat waves, showering is must and also tubewell water. In winter continue adding of tubewell water with KMnO_4 .
(ii) Health and Disease management	Treatment of KMnO_4 @ 10 ppm. Sale out the bigger fishes.	Treatment of KMnO_4 @ 10 ppm. Dump the fishes which were heavily infected	Disinfection with KMnO_4 continues. Sale out all the fishes except, infected ones. Dump the infected fishes in a ditch in the ground.

Annexure 1: Location map of Bhiwani



Annexure 2: Rainfall map

