

State: Madhya Pradesh

Agriculture Contingency Plan for District: Shahdol

1.0 District Agriculture profile			
1.1	Agro-Climatic/Ecological Zone		
	Agro Ecological Sub Region (ICAR)	Central Highlands (Malwa And Bundelkhand), Hot Subhumid (Dry) Eco-Sub region (10.3)	
	Agro-Climatic Zone (Planning Commission)	Eastern Plateau And Hills Region (VII)	
	Agro Climatic Zone (NARP)	North Hill Zone of Chattisgarh (MP-3)	
	List all the districts or part thereof falling under the NARP Zone	Shahdol, Sidhi, Anuppur, Dindori, Mandla and Umaria	
	Geographic coordinates of district headquarters	Latitude	Longitude
		22° 38' to 24° 20' N	80° 28' to 82° 12' E
	Name and address of the concerned ZRS/ ZARS/ RARS/ RRS/ RRTTS	RARS, Rewa	
	Mention the KVK located in the district	Programme Coordinator Krishi Vigyan Kendra, Kalyanpur, Distt. Shahdol- 484 001	
1.2	Rainfall	Normal RF(mm)	Normal Onset
			Normal Cessation
	SW monsoon (June-Sep):	1063.1	2 nd week of June
	NE Monsoon(Oct-Dec):	57	1 st week of October
	Winter (Jan-Feb)	63.8	
	Summer (March-May)	42.1	
	Annual	1226	-

1.3	Land use pattern of the district (latest statistics)	Geographical area	Cultivable area*	Forest area	Land under non-agricultural use	Permanent pastures	Cultivable wasteland	Land under Misc. tree crops and groves	Barren and uncultivable land	Current fallows	Other fallows
	Area ('000 ha)	561.0	231.3	227.8	44.6	6.5	40.8	0.7	9.3	30.8	27.7

* Net sown area + current fallow + old fallow

1.4	Major Soils (common names like red sandy loam deep soils (etc.,))*	Percent (%) of total
	Deep soils	67.3
	Medium deep soils	18.3
	Shallow soils	14.3

Source: NBSS & LUP, Nagpur

1.5	Agricultural land use(2008-09)	Area ('000 ha)	Cropping intensity %
	Net sown area	172.8	117
	Area sown more than once	28.9	
	Gross cropped area	201.7	

(Source : Agriculture Statistics 2009, Directorate of Farmer Welfare and Agriculture Development Madhya Pradesh, Bhopal)

1.6	Irrigation	Area ('000 ha)		
	Net irrigated area	20.8		
	Gross irrigated area	20.8		
	Rainfed area	152.0		
	Sources of Irrigation	Number	Area ('000 ha)	Percentage of total irrigated area
	Canals	76	4.4	21.1
	Tanks	451	2.4	11.5
	Open wells	2470	3.8	18.2
	Bore wells	513	1.3	6.2

	Lift irrigation schemes river	NA	-	
	Micro-irrigation		-	
	Other sources (reservoir)	2865	8.90	42.81
	Total Irrigated Area		20.80	
	Pump sets	10780	-	-
	No. of Tractors	872	-	-
	Groundwater availability and use* (Data source: State/Central Ground water Department /Board)	No. of blocks/ Tehsils	(%) area	Quality of water (specify the problem such as high levels of arsenic, fluoride, saline etc)
	Over exploited	-	-	-
	Critical	-	-	-
	Semi- critical	-	-	-
	Safe	05	-	-
	Wastewater availability and use	-	-	-
	Ground water quality		-	
*over-exploited: groundwater utilization > 100%; critical: 90-100%; semi-critical: 70-90%; safe: <70%				

1.7 Area under major field crops & horticulture etc. (2008-09)

1.7	Major Field Crops cultivated	Area ('000 ha)							
		<i>Kharif</i>			<i>Rabi</i>			Summer	Total
		<i>Irrigated</i>	<i>Rainfed</i>	Total	<i>Irrigated</i>	<i>Rainfed</i>	Total		
	Rice	-		105.1	-			-	105.1
	Kodo Kutki	-		15.0	-			-	15.0
	Maize	-		12.0	-			-	12.0
	Pigeonpea	-		8.2	-			-	8.2
	Blackgram			6.6					6.6
	Sesame			6.3					6.3
	Wheat						20.0	-	20.0
	Mustard						3.2		3.2
	Chickpea						2.9		2.9
	Linseed						2.5		2.5

	Horticulture crops - Fruits	Total area(ha)	Irrigated	Rainfed
	Mango	9.0	-	
	Water Chestnut	22.0	-	
	Goava	20.0	-	
	Others (specify)			
	Horticultural crops - Vegetables	Total area(ha)	Irrigated(ha)	Rainfed
	Okra	124	-	-
	Brinjal	113	-	-
	Potato	610	-	-
	Tomato	162	-	-
	Chilies	47	-	-
	Others (specify)		-	-

(Source : Agriculture Statistics 2009, Directorate of Farmer Welfare and Agriculture Development Madhya Pradesh, Bhopal)

	Medicinal and Aromatic crops -	Total area	Irrigated	Rainfed
	-	NA	-	-
	Others (specify)	-	-	-

	Plantation crops -	Total area	Irrigated	Rainfed
	-	NA	-	-
	Others such as industrial pulpwood crops etc (specify)	-	-	-
	Fodder crops -	Total area	Irrigated	Rainfed
	-	NA	-	-
	Others (specify)	-	-	-
	Total fodder crop area	-	-	-
	Grazing land	-	-	-
	Sericulture etc	-	-	-
	Others (Specify)	-	-	-

1.8	Livestock	Male ('000)	Female ('000)	Total ('000)			
	Non descriptive Cattle (local low yielding)			388.1			
	Crossbred cattle			NA			
	Non descriptive Buffaloes (local low yielding)			NA			
	Graded Buffaloes			110.8			
	Goat			100.7			
	Sheep			8.1			
	Others (Pig ,horse etc)			9.1			
	Commercial dairy farms (Number)	-	-	-			
1.9	Poultry	No. of farms -NA	Total No. of birds ('000)				
	Commercial	-	-				
	Backyard	-	-				
1.10	Fisheries (Data source: Chief Planning Officer)						
	A. Capture						
	i) Marine (Data Source: Fisheries Department)	No. of fishermen	Boats		Nets		Storage facilities (Ice plants etc.)
			Mechanized	Non-mechanized	Mechanized (Trawl nets, Gill nets)	Non-mechanized (Shore Seines, Stake & trap nets)	
		-	-	-	-	-	-
	ii) Inland (Data Source: Fisheries Department)	No. Farmer owned ponds		No. of Reservoirs		No. of village tanks	
		141		14		1379	
	B. Culture						

		Water Spread Area (ha)	Yield (t/ha)	Production ('000 tons)
	i) Brackish water (Data Source: MPEDA/ Fisheries Department)			-
	ii) Fresh water (Data Source: Fisheries Department)	1584-	Village ponds 2000 kg/ha and Reservoirs 70 kg/ha	-
	Others	-	-	-

1.11 Production and Productivity of major crops

1.11	Name of crop	Kharif		Rabi		Summer		Total		Crop residue as fodder ('000
	Rice	83.8	817					83.8	817	
	Kodo Kutki (Minor millets)	6.6	338					6.6	338	
	Maize	10.1	868					10.1	868	
	Pigeonpea	4.0	503					4.0	503	
	Blackgram	2.2	362					2.2	362	
	Wheat			18.5	862			18.5	862	
	Mustard			1.4	370			1.4	370	
	Chickpea			1.3	412			1.3	412	
	Linseed			0.7	263			0.7	263	
	Lentil			0.2	398			0.2	398	
Major Horticultural crops (NA)										
	Mango					-	-	-	-	-
	Aonla					-	-	-	-	-
	Guava					-	-	-	-	-
	Okra-Tomato					-	-	-	-	-
	Brinjal					-	-	-	-	-


(Source : Agriculture Statistics 2009, Directorate of Farmer Welfare and Agriculture Development Madhya Pradesh, Bhopal)

1.12	Sowing window for 5 major field crops (start and end of normal sowing period)	Rice	Kodo	Maize	Wheat	Chickpea
	Kharif- Rainfed	2 nd week of June-3 rd week of July	2 nd week of June – 2 nd week of July	2 nd week of June – 3 rd week of June	-	-
	Kharif-Irrigated	3 rd week of June-3 rd week of July	-	3 rd week of June – 2 nd week of July	-	-
	Rabi- Rainfed	-	-	-	3 rd week of October – 2 nd week of November (up to 10 Nov)	2 nd week of October - 4 th week of October
	Rabi-Irrigated	-	-	-	2 nd week of November-2 nd week of December	1 st week of November – 3 rd week of November


1.13	What is the major contingency the district is prone to? (Tick mark)	Regular	Occasional	None
	Drought	-	√	
	Flood	-	-	√
	Cyclone	-	-	√
	Hail storm	-	√	-
	Heat wave	-	√	-
	Cold wave	-	√	-
	Frost	-	√	-
	Sea water intrusion	-	-	√
	Pests and disease outbreak (specify)	-	√	-



Annexure II



Annexure III



Source: NBSS & LUP, Nagpur

2.0 Strategies for weather related contingencies

2.1 Drought

2.1.1 Rainfed situation

2.1.1 Rainfed situation

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delay by 2 weeks 4 th week of June	Upland unbunded shallow soils	Maize	No change	Follow normal recommended package of practices Timely sowing can be done Dry sowing of paddy Lehi method of sowing in Rice Sowing of Maize by ridge & furrow method	Seed availability, SAU, Beej nigam, NSC, Farmers societies.
		Kodo			
		Kutki			
		Niger			
		Soybean			
	Upland banded shallow (gravelly sandu) soils	Paddy			
		Maize			
		Pigeonpea			
	Lowland banded deep and medium deep soils	Paddy-Chickpea/lentil			
		Paddy-Wheat/ lentil/Mustard			
		Soybean			

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delay by 4 weeks	Upland unbanded shallow soils	Maize	Donot sow maize Prefer alternate crops like	Moisture conservation practices like ridging, conservation furrows,	Seed availability, SAU, Beej nigam, NSC, Farmers

2 nd week of July			Sesame, kodo, kutki, Blackgram, Greengram and Pigeonpea.	dust mulch etc.,	societies.
		Kodo	Kodo- Jawahar Kodo-1, 2, 41, 62, 101, 147, 439, Jawahar-48, Jawahar, 155, JK-106		
		Kutki	Kutki - Jawahar Kutki 1, 2, 8, JK 36		
		Niger	Niger —JNC-6, JNC-1, JNC-9, JVN-1		
		Soybean	Soybean: JS 335, JS 95-60 Or Blackgram – JU-2, JU-3, JU-86, T-9, JBG-623, LBG 684, TAU-1, Berkha, PU-30,35,19 Or Greengram: Pusa vishal, K851, JM721, Jawahar 99 - 37, Hum-1, Hum-2, Tarme-1 L.G.450, T.M.98-50, JM-98-90, PDM 11, 54 and 139		
	Upland banded shallow(gravelly sandu) soils	Paddy	Paddy: JR- 201		
		Maize	Donot sow maize Prefer alternate crops like Sesame, kodo, kutki, Blackgram, Greengram and Pigeonpea.		
		Pigeonpea	Pigeonpea- Pragati ,Jagriti,,Asha ,Nmuber-148,JKM-7,JA-4, Type-21-Pusa-855, ICPL-85063 (Laxmi), JKM-189		
	Lowland banded	Paddy-Chickpea/lentil	Paddy: JR- 201		

	deep and medium deep soils	Paddy-Wheat/ lentil/Mustard		
		Soybean	Soybean: JS 335, JS 95-60	

Condition			Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delay by 6 weeks 4 th week of July	Upland unbunded shallow soils	Maize	Maize intercropping with Caster Donot sow soybean after 10th July Donot sow Maize, Kodo, Kutki, Blackgram and Greegram Prefer alternate crops like kodo, kutki, Sesame and Niger Sesame- TKG -306, TKG-35, JGS-8, JT-21, JT-22, JT-55, PKTS-11, PKTS-12, JT-1 Niger —JNC-6, JNC-1, JNC-9, JVN-1 Kodo- Jawahar Kodo-1, 2, 41, 62, 101, 147, 439, Jawahar-48, Jawahar, 155, JK-106 Kutki - Jawahar Kutki 1, 2, 8, JK 36	Line sowing with seed treatment and balance fertilizer. Line sowing Blade harrowing (Bakhar) for moisture conservation Intercropping of Sesame and niger with Pigeonpea	Seed availability, SAU, Beej nigam, NSC, Farmers societies.
		Kodo			
		Kutki			
		Soybean			
		Niger	Niger —JNC-6, JNC-1, JNC-9, JVN-1		
	Upland banded	Paddy	Prefer to sow alternate crops		

	shallow(gravelly sandu) soils	Maize	like kodo, kutki, Sesame and Niger		
		Pigeonpea			
	Lowland banded deep and medium deep soils	Paddy-Chickpea/lentil	Prefer to sow alternate crops like kodo, kutki, Sesame and Niger (Donot sow soybean after 10th July)		
		Paddy-Wheat/ lentil/Mustard			
		Soybean			

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delay by 8 weeks 2nd week of August	Upland unbanded shallow soils	Maize	Prefer alternate crops, Niger, Castor in kharif and plan for early rabi crops like mustard, linseed ,lentil.	<ol style="list-style-type: none"> 1. Blade harrowing (Bakhar) for moisture conservation 2. Intercropping of Sesame and Niger with Pigeonpea. 3. Moisture conservation by repeat ploughing. 4. Prepration of field for rabi crop 5. Line sowing is preferable 	Source of seed SAU, NSC & SSC For Agronomic Measures the Ongoing scheme like RKVY NREGS etc
		Kodo			
		Kutki			
		Niger			
		Soybean			
	Upland banded shallow(gravelly sandu) soils	Paddy			
		Maize			
		Pigeonpea			
	Lowland banded deep and medium deep soils	Paddy-Chickpea/lentil			
		Paddy-Wheat/ lentil/Mustard			
	Soybean				

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
Early season drought (Normal onset)	Major Farming situation	Normal Crop/cropping system	Crop management	Soil & moisture conservation measures	Remarks on Implementation
Normal onset followed by 15-20 days dry spell after sowing leading to	Upland unbanded shallow soils	Maize	1. Prefer alternate crops like Soybean, Pigeonpea, Greengram and Blackgram on bunds	<ol style="list-style-type: none"> 1. Blade harrowing (Bakhar) for moisture conservation 2. Adopt moisture 	Source of seed SAU, NSC & SSC For Agronomic Measures the
		Kodo			
		Kutki			
		Niger			

poor germination/crop stand etc.		Soybean	2. Weed management by using hand hoe between crop row.	conservation practices. 3. Conservation of excess rain water in high rainfall areas. 4. Mulching. 5. Provide light irrigation through farm pond. 6.Re-sowing,	Ongoing scheme like RKVY NREGS etc
	Upland bunded shallow(gravelly sandu) soils	Paddy	1. Resowing of direct seeded rice 2. Drought resistant varieties of Rice (JR 201),		
		Maize			
		Pigeonpea			
	Lowland bunded deep and medium deep soils	Paddy-Chickpea/lentil	1. Prefer alternate crops like Soybean, Pigeonpea, Greengram and Blackgram on bunds 2. Weed management using hand hoe between crop row. 3. Drought resistant varieties of Rice (JR 201),		
		Paddy-Wheat/ lentil/Mustard			
Soybean					

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Crop management	Soil & moisture conservation measures	Remarks on Implementation
Mid season drought (long dry spell, consecutive 2 weeks rainless (>2.5 mm) period)					
At vegetative stage	Upland unbunded shallow soils	Maize	Life saving irrigation if available Maintain optimum plant population	Provide Supplemental irrigation if available Mulching, Spray of anti-transpirants. Interculture with Dora/Kulpha/Hand hoe in between rows Use uprooted weeds as mulch for moisture	-
		Kodo			
		Kutki			
		Niger			
		Soybean			
	Upland bunded shallow(gravelly sandu) soils	Paddy			
		Maize			
		Pigeonpea			
	Lowland bunded deep and medium deep soils	Paddy-Chickpea/lentil			
		Paddy-Wheat/ lentil/Mustard			
Soybean					

				conservation. Ridges are made after 15-20 lines of crops for the moisture conservation Adopt plant protection measures	
--	--	--	--	--	--

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
Mid season drought (long dry spell)					
At Flowering stage	Upland unbanded shallow soils	Maize	Life saving irrigation if available -	1. Interculture with Dora/Kulpha/H and hoe in between rows . 2. Use of uprooted weeds use as mulch for moisture conservation. 3. Ridges are made after 15-20 lines of crops for the moisture conservation 4. Adopt plant protection measures	-
		Kodo			
		Kutki			
		Niger			
		Soybean			
	Upland banded shallow (gravelly sandu) soils	Paddy			
		Maize			
		Pigeonpea			
	Lowland banded deep and medium deep soils	Paddy-Chickpea/lentil			
		Paddy-Wheat/ lentil/Mustard Soybean			

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Crop management	Rabi Crop planning	Remarks on Implementation
Terminal drought (Early withdrawal of monsoon)					
	Upland unbanded	Maize	1. Life saving irrigation	1. Prefer to sow Lentil,	Source of seed

	shallow soils	Kodo	through sprinkler. 2. Soil moisture conservation by use of mulch. 3. Prefer to sow short duration crop varieties .	Linseed, Chickpea, irrigated and unirrigated wheat 2. Seed treatment with mixture of Thiram (1.5g)+ Carbendazim (1.5g) /kg seed followed by treated with biofertilizers 3. Sowing of small seeded grains mix with FYM and vermicompos 4. Apply light irrigation to Kharif crops for proper grain filling if required and this will helpful in field preparation of <i>Rabi</i> crops	SAU, NSC & SSC For Agronomic Measures the Ongoing scheme like RKVY NREGS etc
		Kutki			
		Niger			
		Soybean			
	Upland banded shallow (gravelly sandu) soils	Paddy			
		Maize			
		Pigeonpea			
	Lowland banded deep and medium deep soils	Paddy-Chickpea/lentil			
		Paddy-Wheat/ lentil/Mustard			
		Soybean			

2.1.2 Irrigated situation

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delayed release of water in canals due to low rainfall	Medium deep to deep soils	Paddy-Wheat/ lentil/Mustard	Green gram-Mustard/ Black gram-Wheat/ Black gram- Chickpea Fallow-Chickpea Medium duration variety of Rice (JRH-4,5,8, MTU 1010, IR-64, PS-3,5,)	Adopt water saving methods like direct seeded rice, SRI Cultivation, Aerobic rice Wheat Prefer short duration low water requirement varieties of wheat. Protective irrigation at CRI stage in wheat. Chickpea should be sown with residual moisture after harvest of soybean or give pre sowing irrigation to chickpea. Maintain optimum plant population	--
		Paddy-Chickpea/lentil			

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Limited release of water in canals due to low rainfall	Medium deep to deep soils	Rice-Wheat	Rice-Chickpea / Green gram-Wheat(Early) Black gram/ Greengram-Wheat Limited irrigation requirement varieties of Wheat (JW 3020, JW 3173, 3269, HW 2004, Sujata) should be sown Limited irrigation requirement varieties of Chickpea (JG 218, 226, 130, 11, 14)	Adopt water saving methods like direct seeding seeded rice, SRI Cultivation, Aerobic rice Blackgram/ Greengram: Adopt <i>in-situ</i> moisture conservation practices at 30DAS Maintain optimum plant population Irrigate at critical stages Conservation tillage Wheat Prefer short duration low water requirement varieties of wheat. Protective irrigation at CRI stage in wheat. Chickpea should be sown with residual moisture after harvest of soybean or give pre sowing irrigation to chickpea	-
		Rice -Chickpea			

Condition			Suggested Contingency measures		
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Non release of water in canals under delayed onset of monsoon in catchment	Medium deep to deep soils	Rice-Wheat	Rice-Chickpea / Green gram-Wheat(Early) Blackgram-Chickpea/ wheat	Blackgram/ Greengram: Adopt <i>in-situ</i> moisture conservation practices at 30DAS Maintain optimum plant population Irrigate at critical stages Conservation tillage Farm bundin Deep ploughing Mulching	-
		Rice -Chickpea			

Condition			Suggested Contingency measures		
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Lack of inflows into tanks due to insufficient /delayed onset of monsoon	Not applicable				

Condition			Suggested Contingency measures		
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Insufficient groundwater recharge due to low rainfall	Not applicable				

2.2 Unusual rains (untimely, unseasonal etc)] (for both rain fed and irrigated situations)

Condition	Suggested contingency measure			
	Vegetative stage	Flowering stage	Crop maturity stage	Post harvest
Continuous high rainfall in a short span leading to water logging				
Rice	<p>Drain the excess water as early as possible</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p> <p>Take up gap filling either with available nursery or by splitting the tillers from the surviving hills</p> <p>Take up suitable plant protection</p> <p>Measures in anticipation of pest & disease out breaks</p>	<p>Drain the excess water as early as possible</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p> <p>Take up suitable plant protection</p> <p>Measures in anticipation of pest & disease out breaks</p>	<p>Drain the excess water as early as possible</p> <p>Take up suitable plant protection measures in anticipation of pest & disease out breaks</p>	<p>Drain out water and spread sheaves loosely in field or field bunds where there is no water stagnation</p> <p>Spray common salt at 5% on panicles to prevent germination and spoilage of straw from moulds</p> <p>Thresh after drying the sheaves properly</p> <p>Ensure proper grain moisture before storing</p>
Maize	<p>Drain the excess water as early as possible</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p> <p>Take up inter cultivation and at optimum soil moisture condition to loosen and aerate the soil and to control weeds</p> <p>Earthing up the crop for anchorage</p> <p>Spray KNO₃ 1 % or water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 at 1% to support nutrition</p> <p>Take up timely control measures for Pink stem borer, sheath blight and Turcicum leaf blight</p>	<p>Drain the excess water as early as possible</p> <p>Apply 20 kg N + 10 kg K /ha after draining excess water</p> <p>Spray KNO₃ 1 % or water soluble fertilizers like 19-19-19, 20-20-20, 21-21-21 at 1% to support nutrition</p> <p>Take up timely control measures for sheath blight and post flowering stalk rots</p>	<p>Drain the excess water as early as possible</p> <p>Allow the crop to dry completely before harvesting</p>	<p>Harvest the cobs after the they are dried up properly.</p> <p>Dry the grain to optimum moisture condition before storing</p>

Pulses & Minor millets	Provide drainage, care should be taken that rain water does not stagnate in the field.	Change care should be taken that rain water does not stagnate in the field.	Care should be taken that rain water does not stagnate in the field.	Produce should be placed under shade. Or protect the produce by tarpaulin kept in T flow
Wheat	Care should be taken that rain water does not stagnate in the field and not allow to top drashing of nitrogenous fertilizers.	Care should be taken that rain water does not stagnate in the field and not allow to top drashing of nitrogenous fertilizers.	Proper drainage should be provided and adopt all plant protection measures	-
Chickpea	Care should be taken that rain water does not stagnate in the field and not allow to top drashing of nitrogenous fertilizers.	Care should be taken that rain water does not stagnate in the field and not allow to top drashing of nitrogenous fertilizers.	Proper drainage should be provided and adopt all plant protection measures	
Heavy rainfall with high speed wind in a short span	Not applicable			
Out break of pests and diseases due to unseasonal rains				
Rice	Spraying of Monocrotophos 36 EC 14 ml or Cypermethrin 10 EC 6 ml per 10 liter of water against stem borer	Spraying of Monocrotophos 36 EC 14 ml or Cypermethrin 10 EC 6 ml per 10 liter of water against stem borer	Removal and destruction of infected panicles due to Loose smut	-
Maize	Plant protection measures for stem borer, army worm. Control stem borer. For control of leaf blight spray Mancozeb @ 2.5g/l.	Plant protection measures for Rust, TLB. Control cob worm and rust PP measures for Stalk rot/rust//TLB by spraying Hexaconazole @ 0.1 %	Plant protection measures for Rust / TLB/Leaf spot in Maize	-
Soybean	Carry out critical survey of fields for insect and disease attack in crops	Carry out critical survey of fields for insect and disease attack in crops	Carry out critical survey of fields for insect and disease attack in crops	-
Wheat	Spray 0.2 % mancozeb 76% WP against wheat rust.	Spray 0.2 % mancozeb 76% WP against wheat rust.	Carry out critical survey of fields for disease attack in crops	
Chickpea	Spray triazophos 40 % EC @ 1-1.5 l/ha in chickpea against pest incidence. · “T” shaped pegs placed in late sown chickpea field for	Spray triazophos 40 % EC @ 1-1.5 l/ha in chickpea against pest incidence. “T” shaped pegs placed in late sown chickpea field for	Spray triazophos 40 % EC @ 1-1.5 l/ha in chickpea against pest incidence. Carry out critical survey of fields for insect and disease attack in	-

	biological control of pod borer and for chemical control spraying of Quinalphos 25 EC or Chloropyriphos 20 EC C or Methyl Parathion 50 EC @ 600 ml dissolve in 500 L of water should be used. Dusting of Fenvalerate 0.4% or Endosulphan 4% 15-20 kg or Quinolphas 1.5 WP 20-25 per hectare with duster.	biological control of pod borer and for chemical control spraying of Quinalphos 25 EC or Chloropyriphos 20 EC C or Methyl Parathion 50 EC @ 600 ml dissolve in 500 L of water should be used. Dusting of Fenvalerate 0.4% or Endosulphan 4% 15-20 kg or Quinolphas 1.5 WP 20-25 per hectare with duster.	crops	
Horticulture				
Tomato	-	Use of Bird perches @ 50/ha. Spray of Endosulfan @ 1.0 Lit /ha.against Fuit borer management at ETL	Spray of Endosulfan @ 1.0 Lit /ha.against Fuit borer management	-
Brinjal	-	Use of Bird perches @ 50/ha. Spray of Endosulfan @ 1.0 Lit /ha.against Fuit & shoot borer management at ETL	Spray of Endosulfan @ 1.0 Lit /ha.against Fuit borer management	-
Chilli	-	Management of Chilli Thrips Use of Imidacloprid @ 3ml/10 lit. of water	Management of Chilli Thrips Use of Imidacloprid @ 3ml/10 lit. of water	-
Cauliflower	-	Management of DBM , Aphids Use of Imidacloprid @ 3ml/10 lit. of water	Management of DBM , Aphids Use of Imidacloprid @ 3ml/10 lit. of water	-

2.3 Floods

Condition	Suggested contingency measure ^o			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Transient water logging/ partial inundation				
Continuous submergence for more than 2 days	Not applicable			
Sea water intrusion				

2.4 Extreme events: Heat wave / Cold wave/Frost/ Hailstorm /Cyclone

Extreme event type	Suggested contingency measure ^r			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Heat Wave	-	-	-	-
Rice	Light and repeated irrigation at the appearance of hair line cracks in soil surface, Correct iron deficiency with 0.5% iron sulphate spray.	Repeated irrigation at the appearance of hair line cracks in soil surface, pounding of water for 15 days after transplanting to check Fe deficiency and for crop establishment.	Repeated irrigation at the appearance of hairline cracks in soil surface	Harvest crop at physiological maturity
Maize, Pigeonpea and Blackgram	Protect the crop with the help of light irrigation, wind breaks are necessary where cold and heat wave in regular	Protect the crop with the help of light irrigation; wind breaks are necessary where cold and heat wave in regular	Protect the crop with the help of light irrigation; wind breaks are necessary where cold and heat wave in regular	Protect the crop with the help of light irrigation
Horticulture	-	-	-	-
Mango , Guava	Protect the crop with the help of light irrigation, wind breaks are necessary where cold and heat wave in regular	Protect the crop with the help of light irrigation; wind breaks are necessary where cold and heat wave in regular	Protect the crop with the help of light irrigation; wind breaks are necessary where cold and heat wave in regular	Harvest at physiological maturity
Cold wave	-	-	-	-
Chick pea Wheat	Light irrigation Smoke generation at night time to rise temperature	Light irrigation Smoke generation at night time to rise temperature	Light irrigation Smoke generation at night time to rise temperature	Harvest at physiological maturity
Frost				
Chickpea, Lentil, Pigeonpea	Give light irrigation, Smoke generation at night time to rise temperature wind breaks are necessary where cold and heat wave in regular	Protect the crop with the help of light irrigation; Smoke generation at night time to rise temperature wind breaks are necessary where cold and heat wave in regular	Protect the crop with the help of light irrigation, Smoke generation at night time to rise temperature wind breaks are necessary	Harvest at physiological maturity -
Horticulture				

Tomato	Delay or late raising of Nursery		Protect the crop with the help of light irrigation, Smoke generation at night time to rise temperature wind breaks are necessary	-
Potato	Cold Tolerant Variety is grown i.e. Pusa Sheetal of Tomato			-
Chilli, Dhania Methi, Cauliflower	-	-		-
Hailstorm	-	-	-	-
Wheat, chickpea	Re-sowing in case of severe damage	Light and frequent irrigation.	<ul style="list-style-type: none"> • Apply 10% additional nitrogen • Light and frequent irrigation 	Timely harvesting and shifting of produce to safer place in case of early forewarning
Mango , Guava- fruit crops	Not applicable	Prune damaged branches and twigs and apply Bordeaux paste 1% to avoid fungal infections	Prune damaged branches and twigs and apply Bordeaux paste 1% to avoid fungal infections Apply hormonal spray NAA 20 ppm + 1 % urea to prevent flower board	Immediate harvesting, grading and marketing of produce
Cyclone	Not applicable			

2.5 Contingent strategies for Livestock, Poultry & Fisheries

2.5.1 Livestock

	Suggested contingency measures		
	Before the event	During the event	After the event
Drought			
Feed and fodder availability	<p>As the district is occasionally prone to drought the following practices may be implemented to prevent fodder shortage problem</p> <p>Sowing of cereals (fodder varieties of Sorghum/ Bajra) and leguminous crops (Lucerne, Berseem, Horse gram, Cowpea) during North-East monsoon under dry land system for fodder production.</p> <p>Collection of soybean and chick pea stover for</p>	<p>Harvest and use biomass of dried up crops (Rice, wheat, Maize, Soybean, Black gram, Green gram, chick pea etc.,) material as fodder</p> <p>Harvest all the top fodder available (Subabul, Glyricidia, Pipol, Prosopis etc) and feed the LS during drought</p> <p>Concentrate ingredients such as Grains,</p>	<p>Encourage progressive farmers to grow multi cut fodder crops of sorghum/bajra/maize with input subsidy</p> <p>Supply of quality stem cuttings of Hybrid napier (CO1), paragrass, guinea grass etc., well before monsoon</p>

	<p>use as feed supplement during drought</p> <p>Preserving the green maize fodder as silage</p> <p>Encourage fodder production with Bajra – stylo-Bajra on rotation basis and also to cultivate short-term fodder crops like sunhemp</p>	<p>brans, chunnies & oilseed cakes, low grade grains etc. unfit for human consumption should be procured from Govt. Godowns for feeding as supplement for high productive animals during drought</p> <p>Promotion of Horse gram as contingent crop and harvesting it at vegetative stage as fodder</p> <p>Continuous supplementation of minerals and vitamin to prevent infertility.</p> <p>Encourage mixing available kitchen waste with dry fodder while feeding to the milch animals</p>	<p>Encourage growing fodder crops like Berseem in winter and Juar in summer season</p> <p>Flushing the stock to recoup</p> <p>Replenish the feed and fodder banks</p>
Drinking water	<p>Adopt various water conservation methods at village level to improve the ground water level for adequate water supply.</p> <p>Identification of water resources</p> <p>De-silting of ponds</p> <p>Rain water harvesting and create water bodies/watering points (when water is scarce use only as drinking water for animals)</p> <p>Construction of drinking water tanks in herding places/village junctions/relief camp locations</p> <p>Community drinking water trough can be arranged in sandies /community grazing areas</p>	<p>Adequate supply of drinking water.</p> <p>Restrict wallowing of animals in water bodies/resources; Add alum in stagnated water bodies</p>	<p>Watershed management practices shall be promoted to conserve the rainwater. Bleach (0.1%) drinking water / water sources</p> <p>Provide clean drinking water</p>
Health and diseases management	<p>Procure and stock emergency medicines and vaccines for important endemic diseases of the</p>	<p>Carryout deworming to all animals entering</p>	<p>Keep close surveillance on</p>

	<p>area</p> <p>All the stock must be immunized for endemic diseases of the area</p> <p>Surveillance and disease monitoring network to be established at Joint Director (Animal Husbandry) office in the district</p> <p>Adequate refreshment training on draught management to be given to VAS, Jr.VAS, LI with regard to health & management measures</p> <p>Procure and stock multivitamins & area specific mineral mixture</p>	<p>into relief camps</p> <p>Identification and quarantine of sick animals</p> <p>Constitution of Rapid Action Veterinary Force</p> <p>Performing ring vaccination (8 km radius) in case of any outbreak</p> <p>Restricting movement of livestock in case of any epidemic</p> <p>Tick control measures be undertaken to prevent tick borne diseases in animals</p> <p>Rescue of sick and injured animals and their treatment</p> <p>Organize with community, daily lifting of dung from relief camps</p>	<p>disease outbreak.</p> <p>Undertake the vaccination depending on need</p> <p>Keep the animal houses clean and spray disinfectants Farmers should be advised to breed their milch animals during July-September so that the peak milk production does not coincide with mid summer</p>
Floods	NA		
Cyclone	NA		
Heat wave and cold wave			
Heat wave	<p>i) Plantation around the shed</p> <p>ii) H₂O sprinklers / foggers in the shed</p> <p>iii) Application of white reflector paint on the roof</p> <p>iv) Thatched sheds should be provided as a shelter to animal to minimize heat stress</p>	<p>Allow the animals early in the morning or late in the evening for grazing during heat waves</p> <p>Feed green fodder/silage / concentrates during day time and roughages / hay during night time in case of heat waves</p> <p>Put on the foggers / sprinklers /fans during heat waves in case of high yielders (Jersey/HF crosses)</p> <p>In severe cases, vitamin 'C' and electrolytes should be added in H₂O during heat waves.</p>	<p>Feed the animals as per routine schedule</p> <p>Allow the animals for grazing (normal timings)</p>
Cold wave	<p>Covering all the wire meshed walls / open area with gunny bags/ polyethylene sheets (with a mechanism for lifting during the day time and</p>	<p>Allow for grazing between 10AM to 3PM during cold waves</p> <p>Add 25-50 ml of edible oil in concentrates and</p>	<p>Feed the animals as per routine schedule</p> <p>Allow the animals for grazing</p>

	putting down during night time)	fed to the animal during cold waves Apply / sprinkle lime powder in the animal shed during cold waves to neutralize ammonia accumulation	(normal timings)
Insurance	Encouraging insurance of livestock	Listing out the details of the dead animals	Submission for insurance claim and availing insurance benefit Purchase of new productive animals

2.5.2 Poultry

	Suggested contingency measures		
	Before the event	During the event	After the event
Drought			
Shortage of feed ingredients	Storing of house hold grain like maize, broken rice etc, in to use as feed in case of severe drought	Supplementation only for productive birds with house hold grain Supplementation of shell grit (calcium) for laying birds Culling of weak birds	Supplementation to all survived birds
Drinking water		Use water sanitizers or offer cool hygienic drinking water	
Health and disease management	Culling of sick birds. De-worming and vaccination against RD and IBD	Mixing of Vit. A,D,E, K and B-complex including vit C in drinking water (5ml in one litre water)	Hygienic and sanitation of poultry house Disposal of dead birds by burning / burying with lime powder in pit
Floods	NA		
Cyclone	NA		
Heat wave and cold wave			

Shelter/environment management	Heat wave: Provision of proper shelter with good ventilation	In severe cases, foggers/water sprinklers/wetting of hanged gunny bags should be arranged Don't allow for scavenging during mid day	Routine practices are followed
	Cold wave: Provision of proper shelter Arrangement for brooding Assure supply of continuous electricity	Close all openings with polythene sheets In severe cases, arrange heaters Don't allow for scavenging during early morning and late evening	Routine practices are followed
Health and disease management	De-worming and vaccination against RD and fowl pox	Supplementation of house hold grain Provide cool and clean drinking water with electrolytes and vit. C In hot summer, add anti-stress probiotics in drinking water or feed	Routine practices are followed

2.5.3 Fisheries/ Aquaculture

	Suggested contingency measures		
	Before the event	During the event	After the event
Drought			
Shallow water in ponds due to insufficient rains/inflow	<ol style="list-style-type: none"> 1. Restricted release of water from reservoir. 2. Supplementary water harvest structures like pond and tanks have to be developed. 3. Renovation and maintenance of existing water harvest structures 	<ol style="list-style-type: none"> 1. Restrict lifting of water for irrigation purpose of crops 2. Catch the stock, market the produce to reduce the density of population in ponds. 	<ol style="list-style-type: none"> 1. Excavate the ponds to increase the depth. 2. Try to release water into the pond if it rains in off-season
Impact of heat & salt load build up in ponds / change in water quality	<ol style="list-style-type: none"> 1. Prepare to release water into the habitat 	<ol style="list-style-type: none"> 1. Mixing of water from the water harvest structure like ponds and tanks into the fish habitat. 	<ol style="list-style-type: none"> 1. Monitoring the water quality and health of aquatic organisms
Floods	NA		
Cyclone	NA		
Heat wave and cold wave			
Management of pond	Good water quality to be maintained, Water	Recirculation of water and pruning	Water treatment with lime

environment	depth to be maintained		
Health and diseases management	Prophylactic measures to be taken	Maintain good quality water in ponds	Treatment of pond water with lime and medicines